

Trace evaluation of matrix determinants and inversion of 4×4 matrices in terms of Dirac covariants

F. Kleefeld and M. Dillig

Institute for Theoretical Physics III, University of Erlangen–Nürnberg,
Staudtstr. 7, D-91051 Erlangen, Germany

e-mail: kleefeld@theorie3.physik.uni-erlangen.de / mdillig@theorie3.physik.uni-erlangen.de¹



Abstract. In the following short paper we list some useful results concerning determinants and inverses of matrices. First we show, how to calculate determinants of $d \times d$ matrices, if their traces are known. As a next step 4×4 matrices are expressed in terms of Dirac covariants. The third step is the calculation of the corresponding inverse matrices in terms of Dirac covariants.

1 Calculation of matrix determinants and permanents in terms of their traces

We denote the determinant of a quadratic $d \times d$ matrix A by:

$$\det A = \begin{vmatrix} A_{11} & \cdots & A_{1d} \\ \vdots & & \vdots \\ A_{d1} & \cdots & A_{dd} \end{vmatrix} \quad (1)$$

The calculation of the determinant of A for different dimensions d is straight forward:

$$\begin{aligned} d = 1 & \Rightarrow \det A = A_{11} \\ d = 2 & \Rightarrow \det A = A_{11}A_{22} - A_{12}A_{21} \\ d = 3 & \Rightarrow \det A = A_{11}A_{22}A_{33} + A_{12}A_{23}A_{31} + A_{13}A_{21}A_{32} - \\ & - A_{31}A_{22}A_{13} - A_{32}A_{23}A_{11} - A_{33}A_{21}A_{12} \end{aligned}$$

¹ Preprint-No. FAU-TP3-97/4

$$\dots \tag{2}$$

In a more unfamiliar way the determinants can be given by:

$$\begin{aligned}
d = 1 \quad \Rightarrow \quad \det A &= \frac{1}{1!} \sum_{i=1}^1 \left\| A_{ii} \right\| = \frac{1}{1!} \sum_{i=1}^1 A_{ii} \\
d = 2 \quad \Rightarrow \quad \det A &= \frac{1}{2!} \sum_{i,j=1}^2 \left\| \begin{array}{cc} A_{ii} & A_{ij} \\ A_{ji} & A_{jj} \end{array} \right\| = \frac{1}{2!} \sum_{i,j=1}^2 (A_{ii}A_{jj} - A_{ij}A_{ji}) \\
d = 3 \quad \Rightarrow \quad \det A &= \frac{1}{3!} \sum_{i,j,k=1}^3 \left\| \begin{array}{ccc} A_{ii} & A_{ij} & A_{ik} \\ A_{ji} & A_{jj} & A_{jk} \\ A_{ki} & A_{kj} & A_{kk} \end{array} \right\| = \\
&= \frac{1}{3!} \sum_{i,j,k=1}^3 (A_{ii}A_{jj}A_{kk} + A_{ij}A_{jk}A_{ki} + A_{ik}A_{ji}A_{kj} - \\
&\quad - A_{ki}A_{jj}A_{ik} - A_{kj}A_{jk}A_{ii} - A_{kk}A_{ji}A_{ij}) \\
d = 4 \quad \Rightarrow \quad \det A &= \frac{1}{4!} \sum_{i,j,k,\ell=1}^4 \left\| \begin{array}{cccc} A_{ii} & A_{ij} & A_{ik} & A_{i\ell} \\ A_{ji} & A_{jj} & A_{jk} & A_{j\ell} \\ A_{ki} & A_{kj} & A_{kk} & A_{k\ell} \\ A_{\ell i} & A_{\ell j} & A_{\ell k} & A_{\ell\ell} \end{array} \right\| = \\
&= \frac{1}{4!} \sum_{i,j,k,\ell=1}^4 (A_{ii}A_{jj}A_{kk}A_{\ell\ell} \pm \dots) \\
d = 5 \quad \Rightarrow \quad \det A &= \frac{1}{5!} \sum_{i,j,k,\ell,m=1}^5 \left\| \begin{array}{ccccc} A_{ii} & A_{ij} & A_{ik} & A_{i\ell} & A_{im} \\ A_{ji} & A_{jj} & A_{jk} & A_{j\ell} & A_{jm} \\ A_{ki} & A_{kj} & A_{kk} & A_{k\ell} & A_{km} \\ A_{\ell i} & A_{\ell j} & A_{\ell k} & A_{\ell\ell} & A_{\ell m} \\ A_{mi} & A_{mj} & A_{mk} & A_{m\ell} & A_{mm} \end{array} \right\| = \\
&= \frac{1}{5!} \sum_{i,j,k,\ell,m=1}^5 (A_{ii}A_{jj}A_{kk}A_{\ell\ell}A_{mm} \pm \dots) \\
\dots & \tag{3}
\end{aligned}$$

From these expansions it is easy to read off the following expressions which evaluate $\det A$ in terms of the $\text{tr}A$:

$$\begin{aligned}
d = 1 \quad \Rightarrow \quad \det A &= \frac{1}{1!} \text{tr}A \\
d = 2 \quad \Rightarrow \quad \det A &= \frac{1}{2!} \{(\text{tr}A)^2 - \text{tr}A^2\} \\
d = 3 \quad \Rightarrow \quad \det A &= \frac{1}{3!} \{(\text{tr}A)^3 - \text{tr}A \text{tr}A^2 + 2 \text{tr}A^3\}
\end{aligned}$$

$$\begin{aligned}
d = 4 &\Rightarrow \det A = \frac{1}{4!} \{(\operatorname{tr} A)^4 - 6 (\operatorname{tr} A)^2 \operatorname{tr} A^2 + 8 \operatorname{tr} A \operatorname{tr} A^3 + 3 (\operatorname{tr} A^2)^2 - 6 \operatorname{tr} A^4\} \\
d = 5 &\Rightarrow \det A = \frac{1}{5!} \{(\operatorname{tr} A)^5 - 10 (\operatorname{tr} A)^3 \operatorname{tr} A^2 + 20 (\operatorname{tr} A)^2 \operatorname{tr} A^3 - 30 \operatorname{tr} A \operatorname{tr} A^4 - \\
&\quad - 20 \operatorname{tr} A^2 \operatorname{tr} A^3 + 15 \operatorname{tr} A (\operatorname{tr} A^2)^2 + 24 \operatorname{tr} A^5\} \\
&\dots
\end{aligned} \tag{4}$$

For permanents the minus signs in the upper sums just have to be replaced by plus signs. In general we obtain the following formula for arbitrary dimension d :

$$\begin{aligned}
\det A &= \frac{1}{d!} \sum_{1k_1+2k_2+\dots+dk_d=d} (-1)^{d-k_1-\dots-k_d} \frac{d!}{1^{k_1} \dots d^{k_d} k_1! \dots k_d!} (\operatorname{tr} A^1)^{k_1} \dots (\operatorname{tr} A^d)^{k_d} \\
&\quad (k_1, \dots, k_d \in \{0, 1, 2, \dots\})
\end{aligned} \tag{5}$$

For permanents we obviously get:

$$\begin{aligned}
\operatorname{per} A &= \frac{1}{d!} \sum_{1k_1+2k_2+\dots+dk_d=d} \frac{d!}{1^{k_1} \dots d^{k_d} k_1! \dots k_d!} (\operatorname{tr} A^1)^{k_1} \dots (\operatorname{tr} A^d)^{k_d} \\
&\quad (k_1, \dots, k_d \in \{0, 1, 2, \dots\})
\end{aligned} \tag{6}$$

The coefficients $d!/(1^{k_1} \dots d^{k_d} k_1! \dots k_d!)$ are discussed e.g. in [Lud96, p. 44].

2 Expansion of 4×4 matrices in terms of Dirac covariants

As a complete linear independent set of sixteen 4×4 matrices we can use the following so called Dirac covariants ($g^{\mu\nu}$ is the metric tensor):

$$\begin{aligned}
&1_4 \\
&\gamma_5 \\
&\gamma^\mu \quad (\mu = 0, 1, 2, 3), \\
&i\gamma_5 \gamma^\mu \quad (\mu = 0, 1, 2, 3) \\
&\sigma^{\mu\nu} \quad (\mu < \nu)(\mu, \nu = 0, 1, 2, 3)
\end{aligned} \tag{7}$$

$$(\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu := 2g^{\mu\nu} 1_4, \quad \gamma_5 := i\gamma^0 \gamma^1 \gamma^2 \gamma^3, \quad \sigma^{\mu\nu} = -\sigma^{\nu\mu} := \frac{i}{2}(\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu) \text{ [Itz88, p. 692]})$$

Apart from the 4 dimensional unit-matrix 1_4 all Dirac covariants are traceless. All Dirac covariants are trace orthogonal, i.e. if Γ and Γ' are two Dirac covariants we obtain:

$$\operatorname{tr}(\Gamma\Gamma') = 0 \quad \text{for} \quad \Gamma \neq \Gamma' \tag{8}$$

Very useful are the following traces:

$$\begin{aligned}
\text{tr}(1_4 1_4) &= 4 \\
\text{tr}(\gamma_5 \gamma_5) &= 4 \\
\text{tr}(\gamma^\mu \gamma^\nu) &= 4 g^{\mu\nu} \\
\text{tr}(i\gamma_5 \gamma^\mu i\gamma_5 \gamma^\nu) &= 4 g^{\mu\nu} \\
\text{tr}(\sigma^{\mu\nu} \sigma^{\rho\sigma}) &= 4 (g^{\mu\rho} g^{\nu\sigma} - g^{\mu\sigma} g^{\nu\rho})
\end{aligned} \tag{9}$$

Using the notation $\not{a} := a_\mu \gamma^\mu$ (Einstein sum convention!) and the completeness of the Dirac covariants we can expand every 4×4 matrix M in terms of the Dirac covariants:

$$M = A 1_4 + B \gamma_5 + \not{C} + i\gamma_5 \not{D} + \frac{1}{2} E_{\mu\nu} \sigma^{\mu\nu} \quad (E_{\mu\nu} = -E_{\nu\mu}) \tag{10}$$

The coefficients are easily calculated by the trace properties of the Dirac covariants:

$$\begin{aligned}
A &= \frac{1}{4} \text{tr}(M) \\
B &= \frac{1}{4} \text{tr}(M \gamma_5) \\
C^\mu &= \frac{1}{4} \text{tr}(M \gamma^\mu) \\
D^\mu &= \frac{1}{4} \text{tr}(M i\gamma_5 \gamma^\mu) \\
E^{\mu\nu} &= \frac{1}{4} \text{tr}(M \sigma^{\mu\nu})
\end{aligned} \tag{11}$$

Products, commutators and anticommutators of Dirac covariants evaluated and expanded in terms of Dirac covariants are given in appendix A.

3 Inversion of 4×4 matrices in terms of Dirac covariants

3.1 General case

Like in the previous section we consider an arbitrary 4×4 matrix expanded in terms of Dirac covariants:

$$M = A 1_4 + B \gamma_5 + \not{C} + i\gamma_5 \not{D} + \frac{1}{2} E_{\mu\nu} \sigma^{\mu\nu} \quad (E_{\mu\nu} = -E_{\nu\mu}) \tag{12}$$

Now we define the following shorthand notations:

$$E_{\mu\nu}^* := \varepsilon_{\mu\nu\rho\sigma} E^{\rho\sigma} \quad (\text{dual tensor of } E^{\mu\nu})$$

$$\begin{aligned}
C \cdot D &:= C_\mu D^\mu \\
C^2 &:= C_\mu C^\mu \quad , \quad D^2 := D_\mu D^\mu \\
(D \pm iC)^2 &:= (D_\mu \pm iC_\mu)(D^\mu \pm iC^\mu) \\
E^2 &:= E_{\mu\nu} E^{\mu\nu} \\
E^* E &:= E_{\mu\nu}^* E^{\mu\nu} =: EE^*
\end{aligned} \tag{13}$$

and observe the following identities:

$$(D - iC)^2 (D + iC)^2 = (D^2 - C^2)^2 + 4(C \cdot D)^2 = (D^2 + C^2)^2 + 4(C \cdot D)^2 - 4C^2 D^2 \tag{14}$$

$$\left(\frac{1}{2} \sigma_{\mu\nu} E^{\mu\nu} \right)^2 = \frac{1}{2} E^2 + \frac{i}{4} \gamma_5 EE^* \tag{15}$$

$$[\not{C} - i\gamma_5 \not{D}] [\not{C} - i\gamma_5 \not{D}] = C^2 + D^2 + 2\gamma_5 \sigma_{\mu\nu} C^\mu D^\nu \tag{16}$$

$$[\not{C} - i\gamma_5 \not{D}] [\not{C} + i\gamma_5 \not{D}] = C^2 - D^2 - 2i\gamma_5 C \cdot D \tag{17}$$

Using the trace formula (4) for $d = 4$ you can verify that:

$$\begin{aligned}
\det M &= \\
&= (D - iC)^2 (D + iC)^2 + \left[(A - B)^2 - \frac{1}{2} E^2 + \frac{i}{4} EE^* \right] \left[(A + B)^2 - \frac{1}{2} E^2 - \frac{i}{4} EE^* \right] - \\
&- 2 \left\{ (A^2 - B^2 + \frac{1}{2} E^2)(D^2 + C^2) + 4 \left[\frac{i}{2} AE_{\mu\nu}^* - BE_{\mu\nu} \right] C^\mu D^\nu - \right. \\
&- \left. 2 E_{\mu}^{\lambda} E_{\lambda\nu} (C^\mu C^\nu + D^\mu D^\nu) - E^{*\lambda}_{\mu} E_{\lambda\nu} (C^\mu D^\nu - D^\mu C^\nu) \right\}
\end{aligned} \tag{18}$$

It is somewhat involved to derive the following expression for the inverse matrix M^{-1} :

$$\begin{aligned}
M^{-1} = \frac{1}{\det M} \left\{ \right. & \left[(A + B\gamma_5 - \frac{1}{2} \sigma_{\mu\nu} E^{\mu\nu})(A - B\gamma_5) - \right. \\
& - [\not{C} - i\gamma_5 \not{D}] [\not{C} - i\gamma_5 \not{D}] \left. \right] [A - B\gamma_5 - \not{C} - i\gamma_5 \not{D}] - \\
& - (A + B\gamma_5 - \frac{1}{2} \sigma_{\mu\nu} E^{\mu\nu}) \left[\frac{1}{2} E^2 - \frac{i}{4} \gamma_5 EE^* - [\not{C} + i\gamma_5 \not{D}] \frac{1}{2} \sigma_{\alpha\beta} E^{\alpha\beta} \right]
\end{aligned}$$

$$- [\not{C} - i\gamma_5 \not{D}] \frac{1}{2} \sigma_{\mu\nu} E^{\mu\nu} [\not{C} - i\gamma_5 \not{D}] \Big\} \quad (19)$$

For some purposes the following expression is more suitable:

$$\begin{aligned} M^{-1} = \frac{1}{\det M} \Big\{ & (A + B\gamma_5 - \frac{1}{2} \sigma_{\mu\nu} E^{\mu\nu}) \left[(A - B\gamma_5)^2 - \frac{1}{2} E^2 + \frac{i}{4} \gamma_5 E E^* - \right. \\ & - [\not{C} + i\gamma_5 \not{D}] (A + B\gamma_5 - \frac{1}{2} \sigma_{\alpha\beta} E^{\alpha\beta}) \Big] - \\ & - [\not{C} - i\gamma_5 \not{D}] \left[(A + B\gamma_5 + \frac{1}{2} \sigma_{\alpha\beta} E^{\alpha\beta}) [\not{C} - i\gamma_5 \not{D}] - \right. \\ & \left. \left. - [\not{C} - i\gamma_5 \not{D}] [\not{C} + i\gamma_5 \not{D}] \right] \right\} \quad (20) \end{aligned}$$

3.2 The case $E^{\mu\nu} = 0$

In this subsection we take the limit $E^{\mu\nu} = 0$, i.e.:

$$M = A 1_4 + B\gamma_5 + \not{C} + i\gamma_5 \not{D} \quad (21)$$

In this limit we obtain for $\det M$:

$$\begin{aligned} \det M &= \\ &= (A^2 - B^2 - C^2 - D^2)^2 + 4(C \cdot D)^2 - 4C^2 D^2 = \\ &= (A^2 - B^2)^2 - 2(A^2 - B^2)(C^2 + D^2) + (C^2 + D^2)^2 + 4(C \cdot D)^2 - 4C^2 D^2 = \\ &= (A^2 - B^2)^2 - 2(A^2 - B^2)(C^2 + D^2) + (C^2 - D^2)^2 + 4(C \cdot D)^2 = \\ &= (A^2 - B^2)^2 - 2(A^2 - B^2)(C^2 + D^2) + (C - iD)^2 (C + iD)^2 = \\ &= (A^2 - B^2 - 2C^2 - 2D^2)(A^2 - B^2) + (C - iD)^2 (C + iD)^2 \quad (22) \end{aligned}$$

The corresponding inverse matrix M^{-1} is:

$$\begin{aligned} M^{-1} &= \frac{1}{\det M} \left[A^2 - B^2 - [\not{C} - i\gamma_5 \not{D}] [\not{C} - i\gamma_5 \not{D}] \right] [A - B\gamma_5 - \not{C} - i\gamma_5 \not{D}] \\ &= \frac{1}{\det M} \left[A^2 - B^2 - C^2 - D^2 - 2\gamma_5 \sigma_{\mu\nu} C^\mu D^\nu \right] [A - B\gamma_5 - \not{C} - i\gamma_5 \not{D}] \quad (23) \end{aligned}$$

If C^μ is proportional to D^μ , i.e. $C^\mu \sim D^\mu$, we get significant simplifications. We note that:

$$C^\mu \sim D^\mu \Rightarrow 4(C \cdot D)^2 - 4C^2 D^2 = 0 \quad , \quad \sigma_{\mu\nu} C^\mu D^\nu = 0 \quad (24)$$

i.e.

$$C^\mu \sim D^\mu \Rightarrow \det M = (A^2 - B^2 - C^2 - D^2)^2 \quad (25)$$

For the inverse matrix we obtain:

$$\begin{aligned} C^\mu \sim D^\mu \Rightarrow M^{-1} &= \frac{1}{\det M} (A^2 - B^2 - C^2 - D^2) [A - B\gamma_5 - \not{C} - i\gamma_5 \not{D}] = \\ &= \frac{A - B\gamma_5 - \not{C} - i\gamma_5 \not{D}}{A^2 - B^2 - C^2 - D^2} \end{aligned} \quad (26)$$

Some special examples with respect to this case are:

$$\begin{aligned} (A + B\gamma_5 + \not{C})^{-1} &= \frac{A - B\gamma_5 - \not{C}}{A^2 - B^2 - C^2} \\ (A + B\gamma_5 + i\gamma_5 \not{D})^{-1} &= \frac{A - B\gamma_5 - i\gamma_5 \not{D}}{A^2 - B^2 - D^2} \end{aligned} \quad (27)$$

Another interesting situation is the case $A = \pm B$, i.e. we start from the matrix:

$$M = A(1_4 \pm \gamma_5) + \not{C} + i\gamma_5 \not{D} \quad (28)$$

Then we get for $\det M$:

$$\det M = (C - iD)^2 (C + iD)^2 \quad (29)$$

The inverse M^{-1} is given by:

$$M^{-1} = \frac{-1}{\det M} [\not{C} - i\gamma_5 \not{D}] [\not{C} - i\gamma_5 \not{D}] [A(1_4 \mp \gamma_5) - \not{C} - i\gamma_5 \not{D}] \quad (30)$$

3.3 The case $C^\mu = D^\mu = 0$

In this subsection we take the limit $C^\mu = D^\mu = 0$, i.e.:

$$M = A 1_4 + B\gamma_5 + \frac{1}{2} E_{\mu\nu} \sigma^{\mu\nu} \quad (E_{\mu\nu} = -E_{\nu\mu}) \quad (31)$$

In this limit we obtain for $\det M$:

$$\det M = \left[(A - B)^2 - \frac{1}{2} E^2 + \frac{i}{4} EE^* \right] \left[(A + B)^2 - \frac{1}{2} E^2 - \frac{i}{4} EE^* \right] \quad (32)$$

The inverse matrix M^{-1} is given by:

$$\begin{aligned}
M^{-1} &= \frac{1}{\det M} (A + B\gamma_5 - \frac{1}{2} \sigma_{\mu\nu} E^{\mu\nu}) \left[(A - B\gamma_5)^2 - \frac{1}{2} E^2 + \frac{i}{4} \gamma_5 E E^* \right] = \\
&= \frac{1}{\det M} \left[(A - B\gamma_5)^2 - \frac{1}{2} E^2 + \frac{i}{4} \gamma_5 E E^* \right] (A + B\gamma_5 - \frac{1}{2} \sigma_{\mu\nu} E^{\mu\nu}) \quad (33)
\end{aligned}$$

A Products, commutators and anticommutators of Dirac covariants

The totally antisymmetric Levi-Civita tensor is defined by [Itz88, p. 692]:

$$\varepsilon_{\mu\nu\rho\sigma} := \begin{vmatrix} g_{\mu 0} & g_{\mu 1} & g_{\mu 2} & g_{\mu 3} \\ g_{\nu 0} & g_{\nu 1} & g_{\nu 2} & g_{\nu 3} \\ g_{\rho 0} & g_{\rho 1} & g_{\rho 2} & g_{\rho 3} \\ g_{\sigma 0} & g_{\sigma 1} & g_{\sigma 2} & g_{\sigma 3} \end{vmatrix} \quad (34)$$

For the commutator and anticommutator of A and B we write:

$$[A, B] := AB - BA \quad , \quad \{A, B\} := AB + BA \quad (35)$$

Some calculation yields the following expansions of products of Dirac covariants in terms of Dirac covariants ($\varepsilon^{\mu\nu}{}_{\lambda\lambda'} \varepsilon^{\lambda\lambda'\rho\sigma} \stackrel{!}{=} -2 (g^{\mu\rho} g^{\nu\sigma} - g^{\mu\sigma} g^{\nu\rho})$):

$1_4 \cdot 1_4 = 1_4$	$1_4 \cdot \gamma_5 = \gamma_5$	$1_4 \cdot \gamma_\mu = \gamma_\mu$
$\gamma_5 \cdot 1_4 = \gamma_5$	$\gamma_5 \cdot \gamma_5 = 1_4$	$\gamma_5 \cdot \gamma_\mu = -i(i\gamma_5\gamma_\mu)$
$\gamma_\mu \cdot 1_4 = \gamma_\mu$	$\gamma_\mu \cdot \gamma_5 = i(i\gamma_5\gamma_\mu)$	$\gamma_\mu \cdot \gamma_\nu = g_{\mu\nu} 1_4 - i\sigma_{\mu\nu}$
$i\gamma_5\gamma_\mu \cdot 1_4 = i\gamma_5\gamma_\mu$	$i\gamma_5\gamma_\mu \cdot \gamma_5 = -i\gamma_\mu$	$i\gamma_5\gamma_\mu \cdot \gamma_\nu = i g_{\mu\nu} \gamma_5 + \frac{i}{2} \varepsilon_{\mu\nu\rho\sigma} \sigma^{\rho\sigma}$
$\sigma_{\mu\nu} \cdot 1_4 = \sigma_{\mu\nu}$	$\sigma_{\mu\nu} \cdot \gamma_5 = \frac{i}{2} \varepsilon_{\mu\nu\rho\sigma} \sigma^{\rho\sigma}$	$\sigma_{\mu\nu} \cdot \gamma_\rho = i\varepsilon_{\sigma\mu\nu\rho} i\gamma_5\gamma^\sigma +$ $+ \frac{i}{2} \varepsilon_{\mu\nu\lambda\lambda'} \varepsilon^{\lambda\lambda'}{}_{\rho\sigma} \gamma^\sigma$

$1_4 \cdot i\gamma_5 \gamma_\mu = i\gamma_5 \gamma_\mu$	$1_4 \cdot \sigma_{\mu\nu} = \sigma_{\mu\nu}$
$\gamma_5 \cdot i\gamma_5 \gamma_\mu = i\gamma_\mu$	$\gamma_5 \cdot \sigma_{\mu\nu} = \frac{i}{2} \varepsilon_{\mu\nu\rho\sigma} \sigma^{\rho\sigma}$
$\gamma_\mu \cdot i\gamma_5 \gamma_\nu = -ig_{\mu\nu} \gamma_5 - \frac{i}{2} \varepsilon_{\mu\nu\rho\sigma} \sigma^{\rho\sigma}$	$\gamma_\rho \cdot \sigma_{\mu\nu} = i \varepsilon_{\sigma\mu\nu\rho} i\gamma_5 \gamma^\sigma -$ $- \frac{i}{2} \varepsilon_{\mu\nu\lambda\lambda'} \varepsilon^{\lambda\lambda'}_{\rho\sigma} \gamma^\sigma$
$i\gamma_5 \gamma_\mu \cdot i\gamma_5 \gamma_\nu = g_{\mu\nu} 1_4 - i\sigma_{\mu\nu}$	$i\gamma_5 \gamma_\rho \cdot \sigma_{\mu\nu} = -i \varepsilon_{\sigma\mu\nu\rho} \gamma^\sigma -$ $- \frac{i}{2} \varepsilon_{\mu\nu\lambda\lambda'} \varepsilon^{\lambda\lambda'}_{\rho\sigma} i\gamma_5 \gamma^\sigma$
$\sigma_{\mu\nu} \cdot i\gamma_5 \gamma_\rho = -i \varepsilon_{\sigma\mu\nu\rho} \gamma^\sigma +$ $+ \frac{i}{2} \varepsilon_{\mu\nu\lambda\lambda'} \varepsilon^{\lambda\lambda'}_{\rho\sigma} i\gamma_5 \gamma^\sigma$	$\sigma_{\mu\nu} \cdot \sigma_{\rho\sigma} = (g_{\mu\rho} g_{\nu\sigma} - g_{\mu\sigma} g_{\nu\rho}) 1_4 +$ $+ i (\sigma_{\mu\sigma} g_{\nu\rho} - \sigma_{\mu\rho} g_{\nu\sigma} +$ $+ g_{\mu\sigma} \sigma_{\nu\rho} - g_{\mu\rho} \sigma_{\nu\sigma}) +$ $+ i \varepsilon_{\mu\nu\rho\sigma} \gamma_5$

(36)

The commutators of the Dirac covariants are evaluated to be:

$[1_4, 1_4] = 0$	$[1_4, \gamma_5] = 0$	$[1_4, \gamma_\mu] = 0$
$[\gamma_5, 1_4] = 0$	$[\gamma_5, \gamma_5] = 0$	$[\gamma_5, \gamma_\mu] = -2i(i\gamma_5 \gamma_\mu)$
$[\gamma_\mu, 1_4] = 0$	$[\gamma_\mu, \gamma_5] = 2i(i\gamma_5 \gamma_\mu)$	$[\gamma_\mu, \gamma_\nu] = -2i\sigma_{\mu\nu}$
$[i\gamma_5 \gamma_\mu, 1_4] = 0$	$[i\gamma_5 \gamma_\mu, \gamma_5] = -2i\gamma_\mu$	$[i\gamma_5 \gamma_\mu, \gamma_\nu] = 2i g_{\mu\nu} \gamma_5$
$[\sigma_{\mu\nu}, 1_4] = 0$	$[\sigma_{\mu\nu}, \gamma_5] = 0$	$[\sigma_{\rho\sigma}, \gamma_\mu] = 2i (\gamma_\rho g_{\mu\sigma} - \gamma_\sigma g_{\mu\rho})$

$[1_4, i\gamma_5\gamma_\mu] = 0$	$[1_4, \sigma_{\mu\nu}] = 0$
$[\gamma_5, i\gamma_5\gamma_\mu] = 2i\gamma_\mu$	$[\gamma_5, \sigma_{\mu\nu}] = 0$
$[\gamma_\mu, i\gamma_5\gamma_\nu] = -2ig_{\mu\nu}\gamma_5$	$[\gamma_\mu, \sigma_{\rho\sigma}] = 2i(\gamma_\sigma g_{\mu\rho} - \gamma_\rho g_{\mu\sigma})$
$[i\gamma_5\gamma_\mu, i\gamma_5\gamma_\nu] = -2i\sigma_{\mu\nu}$	$[i\gamma_5\gamma_\mu, \sigma_{\rho\sigma}] = 2i(i\gamma_5\gamma_\sigma g_{\mu\rho} - i\gamma_5\gamma_\rho g_{\mu\sigma})$
$[\sigma_{\rho\sigma}, i\gamma_5\gamma_\mu] = 2i(i\gamma_5\gamma_\rho g_{\mu\sigma} - i\gamma_5\gamma_\sigma g_{\mu\rho})$	$[\sigma_{\mu\nu}, \sigma_{\rho\sigma}] = 2i(\sigma_{\mu\sigma}g_{\nu\rho} - \sigma_{\mu\rho}g_{\nu\sigma} + g_{\mu\sigma}\sigma_{\nu\rho} - g_{\mu\rho}\sigma_{\nu\sigma})$

(37)

For the anticommutators of the Dirac covariants we obtain:

$\{1_4, 1_4\} = 2 \cdot 1_4$	$\{1_4, \gamma_5\} = 2 \gamma_5$	$\{1_4, \gamma_\mu\} = 2 \gamma_\mu$
$\{\gamma_5, 1_4\} = 2 \gamma_5$	$\{\gamma_5, \gamma_5\} = 2 \cdot 1_4$	$\{\gamma_5, \gamma_\mu\} = 0$
$\{\gamma_\mu, 1_4\} = 2 \gamma_\mu$	$\{\gamma_\mu, \gamma_5\} = 0$	$\{\gamma_\mu, \gamma_\nu\} = 2 g_{\mu\nu} 1_4$
$\{i\gamma_5\gamma_\mu, 1_4\} = 2i\gamma_5\gamma_\mu$	$\{i\gamma_5\gamma_\mu, \gamma_5\} = 0$	$\{i\gamma_5\gamma_\mu, \gamma_\nu\} = i\varepsilon_{\mu\nu\rho\sigma} \sigma^{\rho\sigma}$
$\{\sigma_{\mu\nu}, 1_4\} = 2 \sigma_{\mu\nu}$	$\{\sigma_{\mu\nu}, \gamma_5\} = i\varepsilon_{\mu\nu\rho\sigma} \sigma^{\rho\sigma}$	$\{\sigma_{\mu\nu}, \gamma_\rho\} = 2i\varepsilon_{\sigma\mu\nu\rho} i\gamma_5\gamma^\sigma$

$\{1_4, i\gamma_5\gamma_\mu\} = 2i\gamma_5\gamma_\mu$	$\{1_4, \sigma_{\mu\nu}\} = 2\sigma_{\mu\nu}$
$\{\gamma_5, i\gamma_5\gamma_\nu\} = 0$	$\{\gamma_5, \sigma_{\mu\nu}\} = i\varepsilon_{\mu\nu\rho\sigma}\sigma^{\rho\sigma}$
$\{\gamma_\mu, i\gamma_5\gamma_\nu\} = -i\varepsilon_{\mu\nu\rho\sigma}\sigma^{\rho\sigma}$	$\{\gamma_\rho, \sigma_{\mu\nu}\} = 2i\varepsilon_{\sigma\mu\nu\rho}i\gamma_5\gamma^\sigma$
$\{i\gamma_5\gamma_\mu, i\gamma_5\gamma_\nu\} = 2g_{\mu\nu}1_4$	$\{i\gamma_5\gamma_\rho, \sigma_{\mu\nu}\} = -2i\varepsilon_{\sigma\mu\nu\rho}\gamma^\sigma$
$\{\sigma_{\mu\nu}, i\gamma_5\gamma_\rho\} = -2i\varepsilon_{\sigma\mu\nu\rho}\gamma^\sigma$	$\{\sigma_{\mu\nu}, \sigma_{\rho\sigma}\} = 2(g_{\mu\rho}g_{\nu\sigma} - g_{\mu\sigma}g_{\nu\rho})1_4 + 2i\varepsilon_{\mu\nu\rho\sigma}\gamma_5$

(38)

References

- [Itz88] C. Itzykson, J.-B. Zuber:
Quantum field theory
 (4th printing 1988)
 ©McGraw–Hill, Inc., 1980
- [Lud96] W. Ludwig, C. Falter:
Symmetries in physics (Group theory applied to physical problems)
 (Second edition)
 ©Springer–Verlag Berlin Heidelberg, 1996